

### Summary:


Public health surveillance data indicates levels of influenza like illness (ILI) are trending above the historical 5 year average for Medina County for ILI emergency department (ED) visits and trending below the historical 5 year average for Influenza hospitalizations. The percentage of ILI emergency department (ED) visits among Medina County residents are above the statewide levels (Figure 1). The geographic distribution of ILI ED visits has decreased with influenza activity found in about half of the county's zip codes (Figure 2). There was 1 influenza-associated hospitalizations and 0 novel influenza infections reported to the Medina County Health Department during week 51. The incidence of influenza associated hospitalizations is below the historical average. (Figure 3). The highest frequency of ILI associated hospitalizations are among adults 45 years of age and older and the highest frequency of ILI ED visits were also among children less than 18 years of age (Figure 4). Over-the-counter medication sales for adults are trending upward and over-the-counter medication sales for children are trending upward.

### Dashboard:


Indicator	Current Week	Previous Week	Trend (# of weeks)	Season to Date
Number of ILI ED visits <sup>1</sup>	55	45	 2	345
Percentage of ILI ED visits <sup>1</sup>	3.8%	3.0%	 2	-
Number of Influenza Hospitalizations <sup>2</sup>	1	0	 1	6
Number of Novel Influenza Infections <sup>2</sup>	0	0	 11	0
Number of Over-the-Counter <sup>3</sup> Medication Sales (Adult)*	2081	1856	 1	21733
Number of Over-the-Counter <sup>3</sup> Medication Sales (Children)*	144	130	 1	1407
Number of Influenza Deaths	0	0	 12	0
Number of Pneumonia Deaths	0	1	 1	4

\*Over-the-counter (OTC) medication sales do not reflect all OTC medication sales within the county. Please see Figures 5 and 6 for specific medications that are being tracked and monitored. All numbers are preliminary and subject to change.\*


**Percentage of ILI ED Visits by Week:**




**Geographic Distribution of Medina County ILI ED Visits:**




## Incidence of Influenza-Associated Hospitalizations among Medina County Residents:




## Age Distribution of Influenza ER Visits and Hospitalizations among Medina County Residents:



## Number of OTC Medication Sales for Adults Sold in Medina County Drug Stores:



## Number of OTC Medication Sales for Children Sold in Medina County Drug Stores:



## State, Regional, and National Data:

### **Ohio Surveillance Data:**

- **ODH lab** has reported **97 positive** influenza tests from specimens sent from sentinel ILINet providers and hospital clinical labs. 2019-2020 influenza season results: **(30) A/pdmH1N1; (12) A/H3N2; (55) Influenza B;** (through 12/21/2019).
- The **National Respiratory and Enteric Virus Surveillance System (NREVSS)** has reported **22,916** influenza RT-PCR tests performed at participating facilities. 2019-2020 influenza season positive results: **(28) A/pdmH1N1; (2) A/H3N2; (254) Flu A Not Subtyped; and (860) Flu B;** (12/21/2019).
- **0 pediatric influenza-associated mortalities** have been reported during the 2019-2020 season (through 12/21/2019).
- No **novel influenza A virus infections** have been reported during the 2019-2020 season (through 12/21/2019).
- Incidence of confirmed **influenza-associated hospitalizations** in 2019-2020 season = **616** (through 12/21/2019).

**HHS Regional Surveillance Data\*:** During week 50 (December 8th – December 14th, 2019), the proportion of outpatient visits for ILI in Region 5 (Ohio is in Region 5) was 2.36%, which is **above** the regional baseline of 1.9%. West Virginia reported Local Activity; Michigan reported Regional Activity; Ohio, Pennsylvania, Kentucky, and Indiana reported Widespread Activity.

**National Surveillance Data\*:** During week 50 (December 8th – December 14th, 2019), the majority of the U.S. reported Moderate or High influenza activity, but some states are still reporting Minimal or Low influenza activity. The proportion of outpatient visits for ILI was 3.9%, which is **above** the national baseline of 2.4%. All 10 HHS regions reported ILI levels at or above their region-specific baseline level. Nationally influenza B/Victoria viruses have been reported more frequently than other influenza viruses this season followed by A(H1N1)pdm09. The predominant virus varies by region. The predominant virus also varies by age group. Nationally, influenza B/Victoria viruses are the most commonly reported influenza viruses among children age 0-4 years (49% of reported viruses) and 5-24 years (59% of reported viruses), while A(H3N2) viruses are the most commonly reported influenza viruses among persons 65 years of age and older (42% of reported viruses). Among adults aged 25-64 years, approximately equal proportions of influenza A(H1N1)pdm09 and B/Victoria viruses (39% and 35%, respectively) have been reported.

**\*National-level and regional-level data are reported one week later than Ohio state-level data**

